

Equational axioms associated with finite automata for fixed point operations in cartesian categories

Zoltán Ésik*

Dept. of Computer Science
University of Szeged
Hungary

February 12, 2015

Abstract

The axioms of iteration theories, or iteration categories, capture the equational properties of fixed point operations in several computationally significant categories. Iteration categories may be axiomatized by the Conway identities and identities associated with finite automata. We show that the Conway identities and the identities associated with the members of a subclass $\mathcal{Q}$ of finite automata is complete for iteration categories iff for every finite simple group G there is an automaton $\mathbf{Q} \in \mathcal{Q}$ such that G is a quotient of a group in the monoid $M(\mathbf{Q})$ of the automaton $\mathbf{Q}$. We also prove a stronger result that concerns identities associated with finite automata with a distinguished initial state.

1 Introduction

Most computer scientists are aware of the fact that just about every aspect of their study is related to fixed points. The importance of fixed points is due to the fact that the semantics of recursion and iteration is usually captured by fixed points of functions, functors, or other constructors. Fixed point operations have been widely used in automata and formal language theory and their generalizations, in the semantics of programming languages, abstract data types, process algebra and concurrency, rewriting, programming logics and verification, complexity theory, and many other fields. The study of the equational properties of the fixed point operations has a long history. Some early references are [1, 7, 9, 10, 17, 18, 20]. A general study of the equational properties of fixed point operations has been provided by *iteration theories*, introduced in 1980 independently in [3] and [10]. Main (but somewhat outdated) references to iteration theories are [4] and [6]. The results obtained in [4, 11, 15, 19] show that iteration theories give a full account of the equational properties of fixed point operations.

An iteration theory is a Lawvere theory [16] equipped with a fixed point (or dagger or iteration) operation, subject to a set of equational axioms. A Lawvere theory is a cartesian category such that each object is the n -fold coproduct of a generator object with itself for some integer $n \geq 0$. In this paper, just as in [5, 13, 19], instead of theories, we will consider the slightly more general cartesian categories and refer to iteration theories as iteration categories.

*Work partially supported by the National Foundation of Hungary for Scientific Research (OTKA), grant no. ANN 110883.

Iteration categories (or rather, theories) were axiomatized in [10] by the ‘Conway identities’ (terminology from [4]) and the ‘commutative identities’. The commutative identities are essentially the same as the (‘vector forms’ of the) identities associated with finite automata in [12]. Since each finite group may be seen as an automaton, there is also an identity associated with each finite group.

In [12], it has been shown that the Conway identities, together with a collection of the group identities associated with the groups in a subclass $\mathcal{G}$ of the finite groups is complete for iteration categories iff every finite simple group is a divisor, i.e., a homomorphic image of a subgroup, of a group in $\mathcal{G}$. This result was derived from the fact that for every finite automaton $\mathbf{Q}$, the identity associated with $\mathbf{Q}$ holds in all Conway theories satisfying the identity associated with every simple group which divides the monoid of $\mathbf{Q}$. In Theorem 3.2, we prove the converse of this fact: if a Conway category satisfies the identity associated with a finite automaton $\mathbf{Q}$, then it satisfies all identities associated with the (simple) group divisors of the monoid of $\mathbf{Q}$. The crucial new part of the argument for this result is contained in the proof of Lemma 4.1 that shows that if the identity associated with an automaton $\mathbf{Q}$ holds in a Conway category, then so does each identity associated with any input extension of $\mathbf{Q}$. Using Theorem 3.2, we conclude in Theorem 5.1 that the Conway identities and the collection of the identities associated with a subclass $\mathcal{Q}$ of finite automata is complete for iteration categories iff for every finite simple group G there is an automaton $\mathbf{Q} \in \mathcal{Q}$ such that G divides the monoid of $\mathbf{Q}$. This theorem strengthens a result proved in [13]. Actually we prove an even stronger result (Theorem 3.4) that concerns identities associated with finite automata with a distinguished initial state.

2 Iteration categories

In any category $\mathcal{C}$, we will denote the composition of morphisms $f : A \rightarrow B$ and $g : B \rightarrow C$ by $g \circ f : A \rightarrow C$. The identity morphism corresponding to object A will be denoted $\mathbf{1}_A$.

We will consider *cartesian categories* $\mathcal{C}$ with explicit products. Thus we assume that for any family of objects $C_i, i \in [n] = \{1, \dots, n\}$, there is a fixed product diagram

$$\pi_i^{C_1 \times \dots \times C_n} : C_1 \times \dots \times C_n \rightarrow C_i, \quad i \in [n]$$

with the usual universal property. When $f_i : A \rightarrow C_i, i \in [n]$ is a family of morphisms, the unique mediating morphism $f : A \rightarrow C_1 \times \dots \times C_n$ with $\pi_i^{C_1 \times \dots \times C_n} \circ f = f_i$ for all $i \in [n]$ will be denoted $\langle f_1, \dots, f_n \rangle$. This morphism is called the *tupling* of the f_i . In particular, when $n = 0$, the empty tuple is the unique morphism $!_A : A \rightarrow T$, where T is the fixed terminal object. We will assume that product is ‘associative on the nose’, so that $A \times (B \times C) = (A \times B) \times C$, $A \times T = A = T \times A$, $\langle \mathbf{1}_A, !_b \rangle = \pi_1^{A \times B}$ etc. for all objects A, B, C . In particular, we will assume that for each object A the projection morphism $\pi_1^A : A \rightarrow A$ is the identity morphism $\mathbf{1}_A$. It follows that $\langle f \rangle = f$ for all $f : A \rightarrow B$.

In the sequel, we will call tuplings of projections *base morphism*. Note that any base morphism

$$\langle \pi_{i_1}^{A_1 \times \dots \times A_n}, \dots, \pi_{i_m}^{A_1 \times \dots \times A_n} \rangle : A_1 \times \dots \times A_n \rightarrow A_{i_1} \times \dots \times A_{i_m}$$

with $i_1, \dots, i_m \in [n]$ corresponds to the function $\rho : [m] \rightarrow [n]$ mapping $j \in [m]$ to $i_j \in [n]$. For example, for each A , the *diagonal morphism* $\Delta_{A^n} = \langle \mathbf{1}_A, \dots, \mathbf{1}_A \rangle : A \rightarrow A^n$ corresponds to the unique function $[n] \rightarrow [1]$. We call a base morphism $A_1 \times \dots \times A_n \rightarrow A_{1\rho} \times \dots \times A_{n\rho}$ corresponding to a permutation $\rho : [n] \rightarrow [n]$ a *base permutation*.

In any cartesian category $\mathcal{C}$, we define

$$f \times g = \langle f \circ \pi_1^{A \times B}, g \circ \pi_2^{A \times B} \rangle : A \times B \rightarrow C \times D$$

for all $f : A \rightarrow C, g : B \rightarrow D$. Clearly, $\times$ is a bifunctor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$.

Suppose that $\mathcal{C}$ is a cartesian category. A *dagger operation* on $\mathcal{C}$ maps a morphism $f : A \times C \rightarrow A$ to a morphism $f^\dagger : C \rightarrow A$.

The *Conway identities* are the *parameter* (1), *double dagger* (2) and *composition identities* (3) given below.

$$(f \circ (\mathbf{1}_A \times g))^\dagger = f^\dagger \circ g, \quad (1)$$

all $f : A \times B \rightarrow A, g : C \rightarrow B$,

$$f^{\dagger\dagger} = (f \circ (\Delta_{A^2} \times \mathbf{1}_C))^\dagger, \quad (2)$$

where $f : A \times A \times C \rightarrow A$.

$$(f \circ \langle g, \pi_2^{A \times C} \rangle)^\dagger = f \circ \langle (g \circ \langle f, \pi_2^{B \times C} \rangle)^\dagger, \pi_2^{B \times C} \rangle, \quad (3)$$

for all $f : B \times C \rightarrow A, g : A \times C \rightarrow B$. Note that the *fixed point identity*

$$f^\dagger = f \circ \langle f^\dagger, \mathbf{1}_C \rangle, \quad f : A \times C \rightarrow A \quad (4)$$

is a particular subcase of the composition identity, and the identity

$$(f \circ \pi_2^{A \times C})^\dagger = f, \quad f : C \rightarrow A \quad (5)$$

is a particular subcase of the fixed point identity (4), while

$$(f \circ \pi_2^{A \times B \times C})^\dagger = f^\dagger \circ \pi_1^{B \times C}, \quad f : A \times B \rightarrow A \quad (6)$$

is an instance of the parameter identity (1). The parameter, double dagger and composition identities are sometimes called the identities of *naturality*, *diagonality* and *dinaturality*.

Definition 2.1 [4] A *Conway category* is a cartesian category equipped with a dagger operation satisfying the Conway identities.

Conway categories satisfy several other non-trivial identities including the *Bekić identity* [1] (called the *pairing identity* in [4])

$$\langle f, g \rangle^\dagger = \langle f^\dagger \circ \langle h^\dagger, \mathbf{1}_C \rangle, h \rangle^\dagger, \quad (7)$$

or its ‘dual form’

$$\langle f, g \rangle^\dagger = \langle k^\dagger, \bar{g}^\dagger \circ \langle k^\dagger, \mathbf{1}_C \rangle \rangle, \quad (8)$$

for all $f : A \times B \times C \rightarrow A$ and $g : A \times B \times C \rightarrow B$, where

$$\begin{aligned} h &= g \circ \langle f^\dagger, \mathbf{1}_{B \times C} \rangle : B \times C \rightarrow B \\ k &= f \circ \langle \pi_1^{A \times C}, \bar{g}^\dagger, \pi_2^{A \times C} \rangle : A \times C \rightarrow A \end{aligned}$$

with

$$\bar{g} = g \circ (\langle \pi_2^{A \times B}, \pi_1^{A \times B} \rangle \times \mathbf{1}_C) : B \times A \times C \rightarrow B.$$

By combining the pairing identity (7) with the identities (5) and (6), we obtain

$$\langle f \circ (\mathbf{1}_A \times !_{B \times C}), g \circ (!_A \times \mathbf{1}_{B \times C}) \rangle^\dagger = \langle f^\dagger, g^\dagger \rangle, \quad (9)$$

for all $f : A \times C \rightarrow A$ and $g : B \times C \rightarrow B$. Also,

$$\pi_1^{A \times B} \circ \langle f \circ (\mathbf{1}_A \times !_{B \times C}), g \rangle^\dagger = f^\dagger, \quad (10)$$

for all $f : A \times C \rightarrow A$ and $g : A \times B \times C \rightarrow B$, and

$$\langle f, g \times !_{B \times C} \rangle^\dagger = \langle f \circ (\langle \mathbf{1}_A, g \rangle \times !_{B \times C}), g \times !_{B \times C} \rangle^\dagger \quad (11)$$

$$= \langle (f \circ (\langle \mathbf{1}_A, g \rangle \times \mathbf{1}_C))^\dagger, g \circ (f \circ (\langle \mathbf{1}_A, g \rangle \times \mathbf{1}_C))^\dagger \rangle \quad (12)$$

for all $f : A \times B \times C \rightarrow A$ and $g : A \rightarrow B$.

We will also make use of the *permutation identity* that holds in all Conway categories:

$$(\pi \circ f \circ (\pi^{-1} \times \mathbf{1}_C))^\dagger = \pi \circ f^\dagger, \quad (13)$$

for all $f : A_1 \times \cdots \times A_n \times C \rightarrow A_1 \times \cdots \times A_n$ and all base permutations

$$\pi : A_1 \times \cdots \times A_n \rightarrow A_{1\pi} \times \cdots \times A_{n\pi}.$$

Note also that, by repeated applications of (7), it follows that

$$f^{\dagger \cdots \dagger} = (f \circ (\Delta_{A^n} \times \mathbf{1}_C))^\dagger \quad (14)$$

for all $f : A^n \times C \rightarrow A$, where dagger appears $n \geq 1$ times on the left-hand side.

A concrete description of the free Conway categories and a characterization of the valid identities of Conway categories together with a PSPACE decision algorithm were given in [2]. In the sequel, we will also make use of the following fact.

Lemma 2.2 *Suppose that $f_i : A^{1+n} \times C \rightarrow A$ for each $i \in [n]$ in a Conway category $\mathcal{C}$. Let*

$$g = \langle f_1 \circ (\langle \pi_1^{A^n}, \mathbf{1}_{A^n} \rangle \times \mathbf{1}_C), \dots, f_n \circ (\langle \pi_n^{A^n}, \mathbf{1}_{A^n} \rangle \times \mathbf{1}_C) \rangle : A^n \times C \rightarrow A^n.$$

Then

$$g^\dagger = \langle f_1^\dagger, \dots, f_n^\dagger \rangle^\dagger.$$

Proof. Define $h : A^{2n} \times C \rightarrow A^n$ by

$$h = \langle f_1 \circ (\pi_1^{A^n} \times \mathbf{1}_{A^n \times C}), \dots, f_n \circ (\pi_n^{A^n} \times \mathbf{1}_{A^n \times C}) \rangle.$$

Then, by the double dagger identity (2) and repeated applications of (9),

$$\begin{aligned} g^\dagger &= (h \circ (\langle \mathbf{1}_{A^n}, \mathbf{1}_{A^n} \rangle \times \mathbf{1}_C))^\dagger \\ &= h^{\dagger\dagger} \\ &= \langle f_1^\dagger, \dots, f_n^\dagger \rangle^\dagger. \quad \square \end{aligned}$$

The Conway identities are quite powerful, for example, a general ‘Kleene theorem’ holds in all Conway categories, and both the soundness and relative completeness of Hoare’s logic can be proved just from the Conway identities, cf. [4]. However, they are not complete for the equational theory of the fixed point operations commonly used in computer science. The

missing ingredient is formulated by the notion of identities associated with finite automata, called automaton identities, or identities associated with finite groups, called group identities. In order to define these identities, we need to introduce some definitions and notation.

Suppose that $\mathbf{Q} = (Q, Z, \cdot)$ is a (*deterministic*) *finite automaton*, where Q is the finite nonempty set of states, Z is the finite nonempty set of input letters, and $\cdot : Q \times Z \rightarrow Q$ is the (right) action of Z on Q . The action is extended in the usual way to a right action of Z^* on Q , where Z^* denotes the free monoid of all words over Z including the empty word ϵ . When $q \in Q$ and $u \in Z^*$, we will usually write just qu for $q \cdot u$.

When $u \in Z^*$, we call the function $Q \rightarrow Q$ defined by $q \mapsto qu$ for all $q \in Q$ the *function induced by u* and denote it by $u^{\mathbf{Q}}$. These functions form a (finite) monoid $M(\mathbf{Q})$ whose multiplication is given by $u^{\mathbf{Q}} \cdot v^{\mathbf{Q}} = (uv)^{\mathbf{Q}} = v^{\mathbf{Q}} \circ u^{\mathbf{Q}}$, for all $u, v \in Z^*$.

Suppose that $\mathbf{Q} = (Q, Z, \cdot)$ and $\mathbf{Q}' = (Q, Z', \cdot)$ are finite automata with the same set of states. We say that $\mathbf{Q}'$ is an *extension of $\mathbf{Q}$* if $Z \subseteq Z'$, $a^{\mathbf{Q}} = a^{\mathbf{Q}'}$ for all $a \in Z$, and if for each letter $b \in Z' \setminus Z$ there is some word $u \in Z^*$ with $b^{\mathbf{Q}'} = u^{\mathbf{Q}}$. Moreover, we say that $\mathbf{Q}'$ is a *restriction of $\mathbf{Q}$* if $Z' \subseteq Z$ and $b^{\mathbf{Q}'} = b^{\mathbf{Q}}$ for all $b \in Z'$. Note that when $\mathbf{Q}$ is a restriction of $\mathbf{Q}'$ this does not imply that $\mathbf{Q}'$ is an extension of $\mathbf{Q}$. Also, when $\mathbf{Q}'$ is an extension of $\mathbf{Q}$ then $M(\mathbf{Q}') = M(\mathbf{Q})$, and if $\mathbf{Q}'$ is a restriction of $\mathbf{Q}$ then $M(\mathbf{Q}')$ is a submonoid of $M(\mathbf{Q})$. Each finite automaton $\mathbf{Q} = (Q, Z, \cdot)$ has an extension $\mathbf{Q}' = (Q, Z', \cdot)$ such that for each $u \in Z^*$ there is some $b \in Z'$ with $b^{\mathbf{Q}'} = u^{\mathbf{Q}}$, and for any two finite automata $\mathbf{Q}, \mathbf{Q}'$ with the same state set Q , if $M(\mathbf{Q}) = M(\mathbf{Q}')$ then, up to a renaming of the input letters, $\mathbf{Q}'$ is a restriction of an extension of $\mathbf{Q}$.

Suppose that $\mathbf{Q} = (Q, Z, \cdot)$ is a finite automaton with $Q = \{q_1, \dots, q_n\}$ and $Z = \{a_1, \dots, a_m\}$, say. Let $\mathcal{C}$ be a Conway category and $f : A^m \times C \rightarrow A$ in $\mathcal{C}$. For each $i \in [n]$, let

$$\rho_i^{\mathbf{Q}, A} = \langle \pi_{ia_1}^{A^n}, \dots, \pi_{ia_m}^{A^n} \rangle : A^n \rightarrow A^m,$$

where we identify each state q_i with the integer i . Moreover, let

$$f^{\mathbf{Q}, A} = f \parallel (\rho_1^{\mathbf{Q}, A}, \dots, \rho_n^{\mathbf{Q}, A}) : A^n \times C \rightarrow A^n$$

be the morphism such that for all $i \in [n]$,

$$\pi_i^{A^n} \circ f^{\mathbf{Q}, A} = f \circ (\rho_i^{\mathbf{Q}, A} \times \mathbf{1}_C) : A^n \times C \rightarrow A.$$

Definition 2.3 We say that the identity $\Gamma(\mathbf{Q})$ associated with $\mathbf{Q}$ holds in a Conway category $\mathcal{C}$ if for all $f : A^m \times C \rightarrow A$,

$$(f^{\mathbf{Q}, A})^\dagger = \Delta_{A^n} \circ (f \circ (\Delta_{A^m} \times \mathbf{1}_C))^\dagger.$$

Suppose that a state $q = q_i$ of $\mathbf{Q}$ is distinguished, so that $(\mathbf{Q}, q)$ is an *initialized finite automaton*. We say that the identity $\Gamma(\mathbf{Q}, q)$ holds in $\mathcal{C}$ if

$$\pi_i^{A^n} \circ (f^{\mathbf{Q}, A})^\dagger = (f \circ (\Delta_{A^m} \times \mathbf{1}_C))^\dagger$$

for all $f : A^m \times C \rightarrow A$.

By the permutation identity which holds in all Conway categories, this definition does not depend on the enumeration of the states of $\mathbf{Q}$. It also does not depend on the enumeration of the input letters, since we may replace f with a morphism $g = f \circ \pi$, where $\pi : A^m \rightarrow A^m$ is a base permutation. It is clear that $\Gamma(\mathbf{Q})$ holds in a Conway category $\mathcal{C}$ iff $\Gamma(\mathbf{Q}, q)$ holds in $\mathcal{C}$ for all states q of $\mathbf{Q}$.

Definition 2.4 [10] An *iteration category* is a Conway category satisfying all identities associated with finite automata, or initialized finite automata.

Only iteration categories generated by a single object were considered in [10], called generalized iterative theories, but the generalization is straightforward. See also [5, 13].

Remark 2.5 The first axiomatization of iteration categories (or rather, theories) involved some of the Conway identities and the commutative identities, cf. [4, 10]. In cartesian categories, the identity associated with the a finite automaton is clearly an instance of the commutative identities. Also, in Conway categories, each instance of the commutative identities is easily seen to be equivalent to the identity associated with an automaton.

An example of an iteration category is the category **Cpo** of complete partial orders and continuous (or monotone) functions, equipped with the (parameterized) least fixed point operation as dagger. It is known that an identity involving the cartesian category operations and dagger holds in **Cpo** iff it holds in iteration categories. For this fact and other completeness results see e.g. [4, 6, 10, 15, 11, 19].

It is instructive to explain the meaning of some of the above identities of iteration categories using functional notation, over the category **Cpo**, say, or an appropriate category of sets with structure and structure preserving maps, equipped with dagger. The meaning of the fixed point identity is, of course, that for all $f : A \times C \rightarrow A$ and $y \in C$, $f^\dagger(y)$ is a (canonical) solution of the fixed point equation $x = f(x, y)$. By the parameter identity (1), if $g : D \rightarrow C$, then for all $z \in D$, the canonical solution of $x = f(x, g(z))$ is $f^\dagger(g(z))$.

The double dagger identity (2) asserts that for $f : A \times A \times C \rightarrow C$, the canonical solution of the equation $x = f(x, x, y)$ may be obtained by first solving $x = f(x, y, z)$ to obtain $f^\dagger(y, z)$, and then solving $x = f^\dagger(x, z)$.

Suppose that **Q** is the two-state automaton with 4 input letters inducing the 4 possible state maps. Then the identity $\Gamma(\mathbf{Q})$ associated with **Q** asserts that for all $f : A^4 \times C \rightarrow A$, both components of the canonical solution of the system

$$\begin{aligned} x &= f(x, y, x, y, z) \\ y &= f(y, x, x, y, z) \end{aligned}$$

are equal to the canonical solution of the single equation $x = f(x, x, x, x, z)$. (Here, the input letters are enumerated so that the first letter induces the identity function, the second induces the swapping of the two states, while the last two letters induce the two constant maps.)

As another example, let **Q** be the automaton with n states and a single letter inducing a cyclic permutation. Then $\Gamma(\mathbf{Q})$ asserts that each component of the canonical solution of the system

$$\begin{aligned} x_1 &= f(x_2, y) \\ &\vdots \\ x_{n-1} &= f(x_n, y) \\ x_n &= f(x_1, y) \end{aligned}$$

for $f : A \times C \rightarrow A$ is equal to the solution of the single equation $x = f(x, y)$. In fact, in Conway categories, $\Gamma(\mathbf{Q})$ is equivalent to the *power identity* $(f^n)^\dagger = f^\dagger$, where $f : A \times C \rightarrow A$, and the powers of f are defined by $f^1 = f$ and $f^n = f \circ \langle f^{n-1}, \pi_2^{A \times C} \rangle$, for all $n > 1$.

3 Main results

We may view each finite monoid M as the finite automaton $(M, M, \cdot)$, where the action is given by monoid multiplication. The identity corresponding to this automaton is called the *(monoid) identity associated with the monoid M* . It is denoted $\Gamma(M)$. When M is a group, $\Gamma(M)$ is called a *group identity*.

When M is a finite monoid, by a *group in M* we shall mean a subsemigroup of M which is a group. Thus, a group in M is not required to contain the identity element of M , its unit element may be any idempotent element of M . However, if M is itself a group, then a group in M is just a subgroup of M . Moreover, we say that a (finite) group G *divides* M if G is a quotient of a group in M .

We recall that a finite group G is *simple* if it is nontrivial and its only nontrivial normal subgroup is G itself.

The following result was proved in [12] (see Theorem 16.1).

Theorem 3.1 *Suppose that $\mathcal{C}$ is a Conway category and $\mathbf{Q}$ is a finite automaton. If $\mathcal{C}$ satisfies the identity associated with each simple group divisor of $M(\mathbf{Q})$, then it satisfies the identity associated with $\mathbf{Q}$.*

Our aim is to prove the following converse of Theorem 3.1.

Theorem 3.2 *If a Conway category satisfies the identity $\Gamma(\mathbf{Q})$ associated with a finite automaton $\mathbf{Q}$, then it satisfies the group identity associated with any simple group divisor of $M(\mathbf{Q})$.*

In our argument, we will make use Lemma 4.1, proved below, and some results from [12]:

Theorem 3.3 [12] *Suppose that M is a finite monoid and $\mathcal{C}$ is a Conway category. Then the following are equivalent.*

- $\mathcal{C}$ satisfies the identity $\Gamma(M)$ associated with M .
- $\mathcal{C}$ satisfies the identity associated with every group in M .
- $\mathcal{C}$ satisfies the identity associated with every group dividing M .
- $\mathcal{C}$ satisfies the identity associated with every simple group dividing M .

Indeed, the fact that the first condition implies the second is covered by Corollary 11.3 of [12], and the fact that the second and fourth conditions are equivalent is proved in Proposition 13.5 of [12]. It then follows that the last three conditions are all equivalent. Finally, the last condition implies the first condition by Theorem 16.1 of [12].

Actually we will prove a stronger version of Theorem 3.2. We say that an initialized finite automaton $(\mathbf{Q}, q)$ is *initially connected* if every state of $\mathbf{Q}$ is accessible from q by some word, i.e., when each state of $\mathbf{Q}$ may be written as qu for some word u .

Theorem 3.4 *If a Conway category satisfies the identity $\Gamma(\mathbf{Q}, q)$ associated with an initialized finite automaton $(\mathbf{Q}, q)$, and if $(\mathbf{Q}, q)$ is initially connected, then it satisfies the group identity associated with every simple group divisor of $M(\mathbf{Q})$.*

4 Proof of the main results

In this section, we provide a proof of Theorem 3.2 and Theorem 3.4.

Lemma 4.1 *If an identity associated with a finite automaton holds in a Conway category, then so does the identity associated with any extension of the automaton.*

Proof. Let $\mathcal{C}$ be a Conway category. Suppose that the identity $\Gamma(\mathbf{Q})$ associated with the automaton $\mathbf{Q}$ holds in $\mathcal{C}$, where $\mathbf{Q} = (Q, Z, \cdot)$. Let $Q = \{q_1, \dots, q_n\}$ and $Z = \{a_1, \dots, a_m\}$, say. Below we will identify state q_i with the integer i .

First we consider the extension of $\mathbf{Q}$ with a letter c inducing the identity function, also treated in [13]. So let $\mathbf{Q}' = (Q, Z \cup \{c\}, \cdot)$ be the extension of $\mathbf{Q}$, where c induces the identity function $Q \rightarrow Q$. Let us enumerate the elements of $Z \cup \{c\}$ as $c, a_1, \dots, a_m$ (we may use any order), and suppose that $f : A^{1+m} \times C \rightarrow A$ in $\mathcal{C}$. First we explain the argument using functional notation, where $\mathcal{C}$ is appropriate. The system of fixed point equations corresponding to $\mathbf{Q}'$ with respect to f is

$$\begin{aligned} x_1 &= f(x_1, x_{1a_1}, \dots, x_{1a_m}, y) \\ &\vdots \\ x_n &= f(x_n, x_{na_1}, \dots, x_{na_m}, y), \end{aligned}$$

where for all appropriate i, j , ia_j is the state entered by $\mathbf{Q}$ or $\mathbf{Q}'$ from state i upon receiving the letter a_j . By Lemma 2.2, this system can be solved in two steps. First, let

$$\begin{aligned} x_1 &= f^\dagger(x_{1a_1}, \dots, x_{1a_m}, y) \\ &\vdots \\ x_n &= f^\dagger(x_{na_1}, \dots, x_{na_m}, y) \end{aligned}$$

and then solve this system, which is just the system of fixed point equations corresponding to $\mathbf{Q}$ with respect to $f^\dagger : A^n \times C \rightarrow A$. Since the identity associated with $\mathbf{Q}$ holds, each component of the solution of this latter system is $h^\dagger(y)$, where $h(x, y) = f^\dagger(x, \dots, x, y)$. But by (14), $h^\dagger(y) = k^\dagger(y)$, where $k(x, y) = f(x, \dots, x, y)$.

More formally, we have

$$\rho_i^{\mathbf{Q}', A} = \langle \pi_i^{A^n}, \rho_i^{\mathbf{Q}, A} \rangle : A^n \rightarrow A^{1+m}$$

for all $i \in [n]$, so that

$$\begin{aligned} f^{\mathbf{Q}', A} &= f \parallel (\rho_1^{\mathbf{Q}', A}, \dots, \rho_n^{\mathbf{Q}', A}) \\ &= \langle f \circ (\langle \pi_1^{A^n}, \rho_1^{\mathbf{Q}, A} \rangle \times \mathbf{1}_C), \dots, f \circ (\langle \pi_n^{A^n}, \rho_n^{\mathbf{Q}, A} \rangle \times \mathbf{1}_C) \rangle \\ &= \langle f \circ (\mathbf{1}_A \times \rho_1^{\mathbf{Q}, A} \times \mathbf{1}_C) \circ (\langle \pi_1^{A^n}, \mathbf{1}_{A^n} \rangle \times \mathbf{1}_C), \dots, \\ &\quad f \circ (\mathbf{1}_A \times \rho_n^{\mathbf{Q}, A} \times \mathbf{1}_C) \circ (\langle \pi_n^{A^n}, \mathbf{1}_{A^n} \rangle \times \mathbf{1}_C) \rangle : A^n \times C \rightarrow A^n. \end{aligned}$$

Thus,

$$(f^{\mathbf{Q}', A})^\dagger = \langle f^\dagger \circ (\rho_1^{\mathbf{Q}, A} \times \mathbf{1}_C), \dots, f^\dagger \circ (\rho_n^{\mathbf{Q}, A} \times \mathbf{1}_C) \rangle^\dagger,$$

by Lemma 2.2 and the parameter identity (1),

$$\begin{aligned} &= ((f^\dagger)^{\mathbf{Q}, A})^\dagger \\ &= \Delta_{A^n} \circ (f^\dagger \circ (\Delta_{A^m} \times \mathbf{1}_C))^\dagger, \end{aligned}$$

since $\Gamma(\mathbf{Q})$ holds in $\mathcal{C}$,

$$= \Delta_{A^n} \circ (f \circ (\Delta_{A^{1+m}} \times \mathbf{1}_C))^\dagger,$$

by (14). This proves that the identity $\Gamma(\mathbf{Q}')$ associated with $\mathbf{Q}'$ holds in $\mathcal{C}$.

Next let $\mathbf{Q}' = (Q, Z \cup \{c\}, \cdot)$ be the extension of $\mathbf{Q}$ with an input letter c inducing the same function as the two-letter word ab , for some $a, b \in Z$. We prove that $\Gamma(\mathbf{Q}')$ holds in $\mathcal{C}$. Let $a = a_m$ and $b = a_{j_0}$, say. Moreover, let $g : A^{m+1} \times C \rightarrow A$, and define:

$$\begin{aligned} f_1 &= g \circ (\pi_1^{A^2} \times \cdots \times \pi_1^{A^2} \times \mathbf{1}_{A^2} \times \mathbf{1}_C) : A^{2m} \times C \rightarrow A \\ f_2 &= \pi_{2(j_0-1)+1}^{A^{2m}} \times !_C : A^{2m} \times C \rightarrow A, \end{aligned}$$

where $\pi_1^{A^2}$ appears $m-1$ times in the definition of f_1 . Let $f = \langle f_1, f_2 \rangle : B^m \times C \rightarrow B$, where $B = A^2$. Since $\Gamma(\mathbf{Q})$ holds in $\mathcal{C}$, we have that each component of

$$(f^{\mathbf{Q}, B})^\dagger : C \rightarrow B^n$$

is equal to

$$(f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger : C \rightarrow B.$$

Let $\alpha = \pi_1^{A^2} \times \cdots \times \pi_1^{A^2} : B^n \rightarrow A^n$. Thus,

$$\alpha \circ (f^{\mathbf{Q}, B})^\dagger = \Delta_{A^n} \circ \pi_1^{A^2} \circ (f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger : C \rightarrow A^n. \quad (15)$$

However, we will prove that the left-hand side of (15) is

$$(g^{\mathbf{Q}', A})^\dagger$$

and the right-hand side of (15) is

$$\Delta_{A^n} \circ (g \circ (\Delta_{A^{m+1}} \times \mathbf{1}_C))^\dagger,$$

so that

$$(g^{\mathbf{Q}', A})^\dagger = \Delta_{A^n} \circ (g \circ (\Delta_{A^{m+1}} \times \mathbf{1}_C))^\dagger.$$

To see informally that this holds, suppose that $\mathcal{C}$ is an appropriate category of sets with structure and structure preserving maps. Consider the system of fixed point equations associated with $\mathbf{Q}$ with respect to f ,

$$\begin{aligned} x_i &= f_1(x_{ia_1}, y_{ia_1}, \dots, x_{ia_m}, y_{ia_m}, z) = g(x_{ia_1}, \dots, x_{ia_m}, y_{ia_m}, z) \\ y_i &= f_2(x_{ia_1}, y_{ia_1}, \dots, x_{ia_m}, y_{ia_m}, z) = x_{ia_{j_0}}, \end{aligned}$$

where $i \in [n]$. Here, x_i and y_i range over A and z ranges over C . The left-hand side of (15) corresponds to the canonical solution of this system for the variables $x_1, \dots, x_n$. Now to solve this system, by the Conway identities, we may substitute $x_{ia_{j_0}}$ for y_i on the right-hand side of each equation, for all $i \in [n]$, so that we obtain a new system of fixed point equations

$$\begin{aligned} x_i &= g(x_{ia_1}, \dots, x_{ia_{m-1}}, x_{ia_m}, x_{ia_m a_{j_0}}, z) \\ y_i &= x_{ia_{j_0}}. \end{aligned}$$

For each i , the right-hand side of the equation for x_i does not contain any of the variables y_j . Thus, we may consider the equations for the x_i separately:

$$x_i = g(x_{ia_1}, \dots, x_{ia_m}, x_{iab}, z)$$

for all $i \in [n]$. This is exactly the system of equations corresponding to the automaton $\mathbf{Q}'$ with respect to g . Thus, the left-hand side of (15) is the canonical solution of the system of equations associated with $\mathbf{Q}'$ with respect to g .

Now $(f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger$ is the canonical solution of the system

$$\begin{aligned} x &= f_1(x, y, \dots, x, y, z) = g(x, \dots, x, y, z) \\ y &= f_2(x, y, \dots, x, y, z) = x, \end{aligned}$$

and the first component of the solution is the canonical solution of the single fixed point equation

$$x = g(x, \dots, x, x, z).$$

Thus, each component of the right-hand side of (15) is this canonical solution.

Formally, for each $i \in [n]$, we have

$$\begin{aligned} \pi_i^{B^n} \circ f^{\mathbf{Q}, B} &= \pi_i^{B^n} \circ (f \parallel (\rho_1^{\mathbf{Q}, B}, \dots, \rho_n^{\mathbf{Q}, B})) \\ &= \langle f_1 \circ (\rho_i^{\mathbf{Q}, B} \times \mathbf{1}_C), f_2 \circ (\rho_i^{\mathbf{Q}, B} \times \mathbf{1}_C) \rangle \\ &= \langle g \circ (\langle \pi_1^{A^2} \circ \pi_{ia_1}^{B^n}, \dots, \pi_1^{A^2} \circ \pi_{ia_m}^{B^n}, \pi_2^{A^2} \circ \pi_{ia_m}^{B^n} \rangle \times \mathbf{1}_C), (\pi_1^{A^2} \circ \pi_{ia_{j_0}}^{B^n}) \times !C \rangle. \end{aligned}$$

It follows using (11) and the permutation identity (13) that

$$(f^{\mathbf{Q}, B})^\dagger = h^\dagger,$$

where $h : B^n \times C \rightarrow B^n$ with

$$\begin{aligned} \pi_i^{B^n} \circ h &= \\ &= \langle g \circ (\langle \pi_1^{A^2} \circ \pi_{ia_1}^{B^n}, \dots, \pi_1^{A^2} \circ \pi_{ia_m}^{B^n}, \pi_2^{A^2} \circ \pi_{ia_m a_{j_0}}^{B^n} \rangle \times \mathbf{1}_C), (\pi_1^{A^2} \circ \pi_{ia_{j_0}}^{B^n}) \times !C \rangle \\ &= \langle g \circ (\langle \pi_1^{A^2} \circ \pi_{ia_1}^{B^n}, \dots, \pi_1^{A^2} \circ \pi_{ia_m}^{B^n}, \pi_2^{A^2} \circ \pi_{iab}^{B^n} \rangle \times \mathbf{1}_C), (\pi_1^{A^2} \circ \pi_{ib}^{B^n}) \times !C \rangle \end{aligned}$$

for all $i \in [n]$. Let $\gamma : A^{2n} \rightarrow A^{2n}$ be the base permutation corresponding to the function $[2n] \rightarrow [2n]$ which maps every integer of the form $2(i-1)+1$ for $i \in [n]$ to i , and every integer of the form $2i$ for $i \in [n]$ to $n+i$. Then for each $i \in [n]$, the i th component of $\bar{h} = \gamma^{-1} \circ h \circ (\gamma \times \mathbf{1}_C) : A^{2n} \times C \rightarrow A^{2n}$ is the morphism

$$g \circ (\langle \pi_{ia_1}^{A^n}, \dots, \pi_{ia_m}^{A^n}, \pi_{iab}^{A^n} \rangle \times !A^n \times \mathbf{1}_C) = g \circ (\rho_i^{\mathbf{Q}', A} \times !A^n \times \mathbf{1}_C) : A^{2n} \times C \rightarrow A,$$

so that the first n components of $\bar{h}$ determine the morphism

$$(g \parallel (\rho_1^{\mathbf{Q}', A}, \dots, \rho_m^{\mathbf{Q}', A}, \rho_{m+1}^{\mathbf{Q}', A})) \circ (\mathbf{1}_{A^n} \times !A^n \times \mathbf{1}_C) = g^{\mathbf{Q}', A} \circ (\mathbf{1}_{A^n} \times !A^n \times \mathbf{1}_C) : A^{2n} \times C \rightarrow A^n.$$

Thus, by (10) and the permutation identity, the tupling of the first n components of $\bar{h}^\dagger$ and hence the morphism on the left-hand side of (15) is exactly $(g^{\mathbf{Q}', A})^\dagger : C \rightarrow A^n$.

Now $f \circ (\Delta_{B^m} \times \mathbf{1}_C) : B \times C \rightarrow B$ is

$$\langle g \circ (\Delta_{A^m} \times \mathbf{1}_A \times \mathbf{1}_C), \pi_1^{A^2} \times !C \rangle,$$

and thus, it follows using (12) that

$$\pi_1^{A^2} \circ (f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger = (g \circ (\Delta_{A^{m+1}} \times \mathbf{1}_C))^\dagger.$$

This proves that the morphism on the right-hand side of (15) is $\Delta_{A^n} \circ (g \circ (\Delta_{A^{m+1}} \times \mathbf{1}_C))^\dagger$.

It follows now that if $\mathbf{Q}'$ is an arbitrary extension of $\mathbf{Q}$, then the identity $\Gamma(\mathbf{Q}')$ holds in $\mathcal{C}$. $\square$

Proposition 4.2 *Suppose that $\mathcal{C}$ is a Conway category and $\mathbf{Q}$ is a finite automaton. If the identity associated with $\mathbf{Q}$ holds in $\mathcal{C}$, then so does the identity associated with $M(\mathbf{Q})$.*

Proof. Suppose that $\mathbf{Q} = (Q, Z, \cdot)$, and let $M = M(\mathbf{Q})$. Consider the transformation monoid [8] (Q, M) viewed as an automaton $\mathbf{Q}' = (Q, M, \cdot)$ in the usual way, so that $q \cdot u^{\mathbf{Q}'} = qu$, for all $q \in Q$ and $u \in Z^*$. Similarly, let $\mathbf{Q}_M = (M, M, \cdot)$ be the automaton corresponding to the monoid M . It is well-known that $\mathbf{Q}_M$ is isomorphic to a subautomaton of a direct power of $\mathbf{Q}'$. If the identity associated with $\mathbf{Q}$ holds in $\mathcal{C}$, then so does the identity associated with $\mathbf{Q}'$, by Lemma 4.1. Since the automaton $\mathbf{Q}_M$ associated with M is isomorphic to a subautomaton of a direct power of $\mathbf{Q}_M$, it follows using Lemma 11.2 and Corollary 12.2 of [12] that the identity $\Gamma(\mathbf{Q}_M)$ associated with $\mathbf{Q}_M$ also holds. But the identity associated with $\mathbf{Q}_M$ is just the identity $\Gamma(M)$ associated with M . $\square$

We now complete the proof of Theorem 3.2. Suppose that $\mathcal{C}$ is a Conway category and $\mathbf{Q}$ is a finite automaton such that the identity associated with $\mathbf{Q}$ holds in $\mathcal{C}$. Then, by the previous proposition, the identity associated with $M(\mathbf{Q})$ holds in $\mathcal{C}$. But, by Theorem 3.3, this implies that the identity associated with every (simple) group dividing $M(\mathbf{Q})$ holds in $\mathcal{C}$.

We now turn to the proof of Theorem 3.4. We need the following fact.

Lemma 4.3 *Suppose that $\mathbf{Q} = (Q, Z, \cdot)$ is a finite automaton, $q \in Q$ and $a \in Z$. If the identity $\Gamma(\mathbf{Q}, q)$ holds in a Conway category, then so does the identity $\Gamma(\mathbf{Q}, qa)$.*

Proof. Let $Q = \{q_1, \dots, q_n\}$ and $Z = \{a_1, \dots, a_m\}$. Without loss of generality, we may assume that $q = q_1$ and $a = a_1$. Let us identify each state q_i with the integer i . Suppose that $\mathcal{C}$ is a Conway category satisfying $\Gamma(\mathbf{Q}, q)$. In order to prove that $\mathcal{C}$ also satisfies $\Gamma(\mathbf{Q}, qa)$, let $g : A^m \times C \rightarrow A$ in $\mathcal{C}$, $B = A^2$, and define $f = \langle f_1, f_2 \rangle : B^m \times C \rightarrow B$ by

$$\begin{aligned} f_1 &= g \circ (\pi_1^{A^2} \times \dots \times \pi_1^{A^2} \times \mathbf{1}_C) \\ f_2 &= \pi_1^{A^{2m}} \times \mathbf{1}_C. \end{aligned}$$

Since the identity associated with $(\mathbf{Q}, q)$ holds, we have

$$\pi_1^{B^n} \circ (f^{\mathbf{Q}, B})^\dagger = (f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger. \quad (16)$$

Let γ denote the base permutation $A^{2n} \rightarrow A^{2n}$ corresponding to the function $[2n] \rightarrow [2n]$ given by $(2j-1) \mapsto j$ and $2j \mapsto n+j$, for all $j \in [n]$. Let $h = \gamma^{-1} \circ (f^{\mathbf{Q}, B} \circ (\gamma \times \mathbf{1}_C))$. It is a routine calculation to verify that

$$h = \langle g^{\mathbf{Q}, A} \circ (\mathbf{1}_{A^n} \times \mathbf{1}_{A^n} \times \mathbf{1}_C), \langle \pi_{1a_1}^{A^n}, \dots, \pi_{na_1}^{A^n} \rangle \times \mathbf{1}_{A^n \times C} \rangle : A^{2n} \times C \rightarrow A^{2n}.$$

Now

$$h^\dagger = \langle (g^{\mathbf{Q}, A})^\dagger, \langle \pi_{1a_1}^{A^n}, \dots, \pi_{na_1}^{A^n} \rangle \circ (g^{\mathbf{Q}, A})^\dagger \rangle : C \rightarrow A^{2n},$$

by (12), and thus by the permutation identity,

$$\pi_1^{B^n} \circ (f^{\mathbf{Q}, A})^\dagger = \langle \pi_1^{A^n} \circ (g^{\mathbf{Q}, A})^\dagger, \pi_{1a_1}^{A^n} \circ (g^{\mathbf{Q}, A})^\dagger \rangle : C \rightarrow A^2.$$

On the other hand, by a similar argument,

$$\begin{aligned} (f \circ (\Delta_{B^m} \times \mathbf{1}_C))^\dagger &= \langle g \circ (\Delta_{A^m} \times \mathbf{1}_{A^m} \times \mathbf{1}_C), \mathbf{1}_A \times \mathbf{1}_A \times \mathbf{1}_C \rangle^\dagger \\ &= \Delta_{A^2} \circ (g \circ (\Delta_{A^m} \times \mathbf{1}_C))^\dagger : C \rightarrow A^2. \end{aligned}$$

Since (16) holds, it follows that $\pi_1^{A^n} \circ (f^{\mathbf{Q}, A})^\dagger = (g \circ (\Delta_{A^m} \times \mathbf{1}_C))^\dagger$. $\square$

By repeated applications of Lemma 4.3 we obtain:

Corollary 4.4 *Suppose that $\mathbf{Q}$ is an automaton and q is a state of $\mathbf{Q}$ and u is an input word of $\mathbf{Q}$. If the identity $\Gamma(\mathbf{Q}, q)$ holds in a Conway category $\mathcal{C}$, then so does the identity $\Gamma(\mathbf{Q}, qu)$. Thus, if $(\mathbf{Q}, q)$ is initially connected, then $\Gamma(\mathbf{Q})$ holds in a Conway category $\mathcal{C}$ iff $\Gamma(\mathbf{Q}, q)$ does.*

Remark 4.5 Our proof of Corollary 4.4 based on Lemma 4.3 uses essentially the same technique as the proof of Lemma 4.3 in [19].

Theorem 3.4 is now an immediate corollary of Theorem 3.2 and Corollary 4.4.

5 Some corollaries and open problems

Combining Theorem 3.1 and Theorem 3.2, we immediately have:

Theorem 5.1 *Suppose that $\mathcal{Q}$ is a collection of finite automata and let $\mathcal{G}$ denote the collection of simple groups dividing the monoid of some automaton in $\mathcal{Q}$. Then a Conway category satisfies the identities associated with the automata in $\mathcal{Q}$ iff it satisfies the identities associated with the groups in $\mathcal{G}$.*

Proof. Suppose that a Conway category $\mathcal{C}$ satisfies the identities $\Gamma(G)$ associated with the groups G in $\mathcal{G}$. Then by Theorem 3.1, $\mathcal{C}$ satisfies all identities $\Gamma(\mathbf{Q})$, where $\mathbf{Q}$ is a finite automaton such that every simple group divisor of $M(\mathbf{Q})$ is in $\mathcal{G}$ and thus all the identities associated with the finite automata in $\mathcal{Q}$.

Suppose now that $\mathcal{C}$ is a Conway category satisfying the identities associated with the finite automata in $\mathcal{Q}$. Then by Theorem 3.2, every identity associated with the groups in $\mathcal{G}$ holds in $\mathcal{C}$. $\square$

Similarly, we have:

Theorem 5.2 *Suppose that $\mathcal{Q}$ is a collection of finite initially connected initialized automata, and let $\mathcal{G}$ denote the collection of simple groups dividing the monoid of some automaton in $\mathcal{Q}$. Then a Conway category satisfies the identities associated with the automata in $\mathcal{Q}$ iff it satisfies the identities associated with the groups in $\mathcal{G}$.*

Corollary 5.3 *Suppose that $\mathcal{Q}$ is a collection of finite automata, or finite initially connected initialized automata. Then the Conway identities together with the identities associated with the (initialized) automata in $\mathcal{Q}$ is complete for iteration categories iff for every finite simple group G there is an automaton $\mathbf{Q} \in \mathcal{Q}$ (or initialized automaton $(\mathbf{Q}, q) \in \mathcal{Q}$) such that G divides the monoid $M(\mathbf{Q})$.*

Remark 5.4 This Corollary strengthens a result proved in [13] using some more sophisticated methods from the structure theory of finite automata. Suppose that $\mathcal{Q}$ is a collection of finite automata such that at least one nontrivial counter is a subautomaton of a restriction of an automaton in $\mathcal{Q}$, i.e., such that there is an integer $n > 1$ and an automaton $\mathbf{Q} \in \mathcal{Q}$ having an n -element subset of the states and an input letter z inducing a cyclic permutation of this subset. It was proved in [13] that the Conway identities and the identities associated with the automata in a collection of finite automata $\mathcal{Q}$ form a complete set of axioms for

iteration categories iff for every finite simple group G there is an automaton $\mathbf{Q} \in \mathcal{Q}$ such that G divides the monoid $M(\mathbf{Q})$.

We note that there is a class $\mathcal{Q}$ of finite automata such that every finite group is isomorphic to the monoid of some automaton in $\mathcal{Q}$ but there is no nontrivial counter that is isomorphic to a subautomaton of a restriction of any automaton in $\mathcal{Q}$.

For each $n \geq 3$, consider the n -state automaton $\mathbf{Q}_n$ with state set $\{q_1, \dots, q_n\}$ with an input letter inducing the cyclic permutation $(q_1 \dots q_n)$ and a letter inducing the transposition (12) , so that the monoid of $\mathbf{Q}_n$ is the symmetric group S_n of degree n . Then the Conway identities together with any infinite subsystem of the identities associated with the initialized automata $(\mathbf{Q}_n, q_1)$ are complete for iteration categories since every finite group may be embedded in any large enough symmetric group. It is shown in [13] that the identity associated with $(\mathbf{Q}_n, q_1)$ may be reduced to

$$(f \circ (\Delta_{A^2} \times \mathbf{1}_C)) \circ (f \circ \langle \pi_1^{A \times C}, (f^\dagger)^{n-2}, \pi_2^{A \times C} \rangle, \pi_2^{A \times C})^\dagger = (f \circ (\Delta_{A^2} \times \mathbf{1}_C))^\dagger, \quad (17)$$

where f is any morphism $A^2 \times C \rightarrow A$. Thus, we obtain a simple proof of the fact that system consisting of the Conway identities and any infinite subset of the identities (17) is complete for iteration categories.

None of the complete systems consisting of the Conway identities and an infinite subcollection of the identities (17) is minimal, since each such system contains a complete proper subsystem. Similarly, no complete axiomatization of iteration theories consisting of the Conway identities and a subcollection of the automaton (or group) identities is minimal.

Open problem Does there exist a minimal complete set of identities for iteration categories?

Given a class $\mathcal{G}$ of finite simple groups (including the trivial groups), closed under division, there is a corresponding ‘variety’ $\mathbf{V}_{\mathcal{G}}$ of Conway categories axiomatized by the Conway identities and the identities $\Gamma(G)$ for $G \in \mathcal{G}$. It contains the subvariety $\mathbf{V}_{\mathcal{G}}^s$ consisting of those categories in $\mathbf{V}_{\mathcal{G}}$ with a single morphism $T \rightarrow A$ for any object A . (Here T is the distinguished terminal object.) These varieties are all different. A concrete description of the ‘free categories’ (or rather, free iteration theories) in these varieties was given in [14]. It is possible that all varieties of Conway categories are of the form $\mathbf{V}_{\mathcal{G}}$ or $\mathbf{V}_{\mathcal{G}}^s$. By the results of this paper, this holds iff every variety of Conway categories may be axiomatized by the Conway identities and a set of automaton identities, and possibly the identity $f = g$, where $f, g : T \rightarrow A$.

Open problem Describe the lattice of all varieties of Conway categories.

6 Conclusion

The equational properties of fixed point operations over cartesian categories are captured by the axioms of iteration categories which can be axiomatized by the Conway identities and identities associated with (deterministic) finite automata, or finite groups. We proved that in conjunction with the Conway identities, the identity associated with a finite automaton implies each identity associated with every finite group that divides the monoid of the automaton. We concluded that a system consisting of the Conway identities and a collection of the identities associated with the members $\mathbf{Q}$ of a subclass $\mathcal{Q}$ of finite automata is complete for iteration categories iff for each finite simple group G there is some automaton $\mathbf{Q} \in \mathcal{Q}$ such that G divides the monoid of $\mathbf{Q}$. This result establishes a connection between the

equational theory of fixed point operations and the Krohn-Rhodes decomposition theorem for finite automata. Since several equational theories of computational interest have a finite axiomatization relatively to iteration categories, our result yields simple axiomatizations of these theories.

References

- [1] H. Bekić, Definable operations in general algebras, and the theory of automata and flowcharts. *Technical Report*, IBM Laboratory, Vienna, 1969. Reprinted in: *Programming Languages and their Definition*. Selected papers edited by C. B. Jones. LNCS 177, Springer-Verlag, 1984, 30–55.
- [2] L. Bernátsky and Z. Ésik, Semantics of flowchart programs and the free Conway theories. *Theoretical Informatics and Applications*, 32(1998), 35–78.
- [3] S.L. Bloom, C.C. Elgot, and J.B. Wright. Solutions of the iteration equation and extensions of the scalar iteration operation. *SIAM Journal of Computing*, 9(1980), 26–45.
- [4] S.L. Bloom and Z. Ésik. *Iteration Theories: The Equational Logic of Iterative Processes*. EATCS Monographs on Theoretical Computer Science. Springer-Verlag, 1993.
- [5] S.L. Bloom and Z. Ésik. Fixed-point operations on ccc’s. Part I. *Theoretical Computer Science*, 155(1996), 1–38.
- [6] S.L. Bloom and Z. Ésik. The equational logic of fixed points, *Theoretical Computer Science*, 179(1997), 1–60.
- [7] J.W. DeBakker and D. Scott. A theory of programs, *Technical Report*, IBM Vienna, 1969.
- [8] S. Eilenberg. *Automata, Languages, and Machines*, vol. B, Academic Press, 1976.
- [9] C.C. Elgot. Monadic computation and iterative algebraic theories. In J.C. Shepherdson, editor, *Logic Colloquium 1973, Studies in Logic*, volume 80. North Holland, Amsterdam, 1975. 175–230.
- [10] Z. Ésik. Identities in iterative and rational theories. *Computational Linguistics and Computer Languages*, 14(1980), 183–207.
- [11] Z. Ésik. Completeness of Park induction. *Theoretical Computer Science*, 177(1997), 217–283.
- [12] Z. Ésik. Group axioms for iteration. *Information and Computation*, 148(1999), 131–180.
- [13] Z. Ésik. Axiomatizing iteration categories. *Acta Cybernetica*, 14(1999), 65–82.
- [14] Z. Ésik. The power of the group axioms for iteration. *International J. Algebra and Computation*, 10(2000), 349–373.
- [15] Z. Ésik and A. Labella. Equational properties of iteration in algebraically complete categories. *Theoretical Computer Science*, 195(1998), 61–89.
- [16] F.W. Lawvere. Functorial semantics of algebraic theories. *Proceedings of the National Academy of Sciences USA*, 50(1963), 869–873.

- [17] D. Niwinski. On fixed-point clones (Extended abstract), *Automata, Languages and Programming, 13th International Colloquium, ICALP86*, LNCS 226, Springer-Verlag, 1986, 464–473.
- [18] G. Plotkin. *Domains*. The Pisa notes. The University of Edinburgh, 1983.
- [19] A. Simpson and G. Plotkin. Complete axioms for categorical fixed-point operators, *15th Annual IEEE Symposium on Logic in Computer Science*, IEEE Computer Society, 2000, 30–41.
- [20] J.B. Wright, J.W. Thatcher, E.G. Wagner and J.A. Goguen. Rational algebraic theories and fixed-point solutions. In *FOCS 1976*, IEEE Computer Society, 1976, 147–158.